

letters

Withdrawal of two generic clopidogrel products in Saudi Arabia for non-bioequivalence

To the Editor: Clopidogrel, a P2Y₁₂ platelet inhibitor, is indicated for the treatment of cardiovascular and cerebrovascular events.¹⁻³ In February 2018, the Saudi Food and Drug Authority (SFDA) announced the withdrawal of two locally approved clopidogrel generic products (Pedovex and Cardlet), secondary to failure in demonstrating bioequivalence to the reference product (Plavix). With this action, the SFDA has reconfirmed that annual post-marketing surveillance is a continuous ongoing process that ensures that all marketed products are up to standards.

Pedovex was withdrawn after retesting in a new bioequivalence study.⁴ The retesting was prompted by several lack of efficacy reports received by the SFDA. In its announcement, the SFDA provided no specific information on how much the withdrawn generics deviated from the reference product. However, the reports of reduced efficacy suggested that these generics failed because of low bioavailability.

The withdrawal of these two local clopidogrel generic products attracted the attention of healthcare professionals in Saudi Arabia because of the vital role of clopidogrel in preventing recurrent stroke and heart ischemia. The action undertaken by the SFDA has also generated concerns among healthcare professionals regarding the quality of some generic products in use in Saudi Arabia. To be approved by regulatory agencies, a generic drug needs to be "bioequivalent" to its brand-name counterpart. The criterion for establishing bioequivalence specifies that the 90% confidence

interval (CI) for the geometric mean ratio (test:reference) for the area under the curve (AUC) and maximum concentration (C_{max}), lie between 80% and 125%.⁵ This criterion is based on the concept that a 20% change in C_{max} /AUC values is clinically acceptable. This range is reasonable for most drugs. However, it might be considered too wide for drugs with a narrow therapeutic index.⁶⁻⁸ Therefore, some regulatory agencies have set more stringent criteria to establish bioequivalence for drugs with a narrow therapeutic index. For example, the European Medicines Agency (EMA) requires that the 90% CI for the AUC lie between 90% and 112%.⁸ Clopidogrel might be considered in this class.^{9,10} Several studies have shown that a modest decrease in the AUC value of the active metabolite of clopidogrel could lead to treatment failure, while a higher AUC value could lead to an increased risk of bleeding.¹¹⁻¹⁵

Designing a study to establish bioequivalence for clopidogrel generic products is complex. As a prodrug, it requires in vivo biological transformation to yield the active moiety. Furthermore, there is high between-subject variability in its pharmacokinetics. In the case of prodrugs, both the U.S. Food and Drug Administration (FDA) and EMA recommend that the parent drug, not the active metabolite, be measured. The old analytical method is based on measuring the active metabolite whereas the new technique is based on the parent drug. It is important to note that the SFDA bioequivalence study was based on the new analytical method (measurement of the parent drug), while the two withdrawn generic products were initially registered using the old analytical method (measurement of the active metabolite). Currently, the SFDA is reviewing all clopidogrel generic products that

were approved based on the old method (measurement of the active metabolite). This is an important step by the SFDA to ensure the quality of clopidogrel generic products because any issues with the absorption of clopidogrel would lead to underexposure and could be associated with severe outcomes such as strokes and myocardial infarctions.

Compliance with Good Manufacturing Process is another issue associated with these products. Once a generic product is approved for use by regulatory agencies, retesting for bioequivalence is usually not mandated. However, there are other measures to confirm the quality of the generic product after approval. Regulatory agencies require companies to perform routine testing, especially for new batch releases. Furthermore, the SFDA has a post-marketing surveillance system that allows healthcare professionals to report issues such as adverse effects and drug quality defects. This system played an important role in the recent events. The SFDA performed the bioequivalence study based on reports from healthcare professionals of a lack of efficacy with the new generics.

Generic medications are gaining acceptance worldwide, primarily because they cost less than their brand-name counterparts. The market for generic drugs is huge. According to IMS Health, in the US alone, unbranded generic drugs accounted for 80% of prescriptions dispensed during the 2013 fiscal year. It is therefore important to maintain the confidence of healthcare providers and patients in generic products. Using generic medications results in savings for patients, payers, and healthcare systems. Many studies have shown that generics can be equal in efficacy to their brand-name counterparts.¹⁶⁻²⁰

Maintaining the confidence of the public and healthcare professionals in generic products requires efforts by regulatory agencies and cooperation from healthcare professionals. An important function of a high-performing healthcare system is ensuring equitable access to essential medical products that are of assured quality, safety, efficacy, and cost-effectiveness.²¹ Post-marketing analyses of generic medications performed by the SFDA could help increase public confidence in these products.

We applaud the actions undertaken by the SFDA in investigating and subsequently withdrawing these generic products of clopidogrel that failed to demonstrate bioequivalence. These actions should increase the vigilance of healthcare professionals toward medication outcomes and encourage the reporting of any unusual efficacy and safety outcomes to the SFDA. We also encourage academia and the commercial sector to perform post-marketing surveillance studies of generic medications. Such initiatives should further help in ensuring the quality of generic products after approval.

Abdullah Saleh Alsultan,^a Hakeam Abdulaziz Hakeam^b

From the ^aDepartment of Pharmacy, King Saud University, Riyadh, Saudi Arabia; ^bPharmacy Services, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

Correspondence: Dr. Abdullah Alsultan, Clinical Pharmacy Department, College of Pharmacy, King Saud University, P. O. Box 2457, Riyadh 11451, Saudi Arabia
T: +96614677485 absultan@ksu.edu.sa
ORCID: <http://orcid.org/0000-0002-7013-8369>

Citation: Alsultan A, Hakeam HA. Withdrawal of two generic clopidogrel products in Saudi Arabia for non-bioequivalence. *Ann Saudi Med* 2018 38(3): 233-234
DOI: 10.5144/0256-4947.2018.233

Received: April 1, 2018

Accepted: April 21, 2018

Published: April 30, 2018

Copyright: Copyright © 2018, Annals of Saudi Medicine, Saudi Arabia. This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND). The details of which can be accessed at <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Funding: None.

REFERENCES

1. Yusuf S, Zhao F, Mehta SR, et al. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med*. 2001;345:494-502.
2. Wang Y, Wang Y, Zhao X, Liu L, Wang D, Wang C, et al. Clopidogrel with aspirin in acute minor stroke or transient ischemic attack. *N Engl J Med*. 2013;369:11-19.
3. Sabatine MS, Cannon CP, Gibson CM, Lopez-Sendon JL, Montalescot G, Theroux P, et al. Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Engl J Med*. 2005;352:1179-1189.
4. SFDA. "SFDA Cancels Registration of Four Products Containing Active Clopidogrel Ingredient". Vol March 18 2018: SFDA; February 2018.
5. Galgatte UC, Jamdade VR, Aute PP, Chaudhari PD. Study on requirements of bioequivalence for registration of pharmaceutical products in USA, Europe and Canada. *Saudi Pharm J*. 2014;22:391-402.
6. Tamargo J, Le Heuzey JY, Mabo P. Narrow therapeutic index drugs: a clinical pharmacological consideration to flecainide. *Eur J Clin Pharmacol*. 2015;71:549-567.
7. Meredith P. Bioequivalence and other unresolved issues in generic drug substitution. *Clin Ther*. 2003;25:2875-2890.
8. Garcia-Arieta A, Gordon J. Bioequivalence requirements in the European Union: critical discussion. *AAPS J*. 2012;14:738-748.
9. Roden DM, Johnson JA, Kimmel SE, Krauss RM, Medina MW, Shuldiner A, et al. Cardiovascular pharmacogenomics. *Circ Res*. 2011;109:807-820.
10. Wang L, McLeod HL, Weinshilboum RM. Genomics and drug response. *N Engl J Med*. 2011;364:1144-1153.
11. Mega JL, Close SL, Wiviott SD, Shen L, Hockett RD, Brandt JT, et al. Cytochrome p-450 polymorphisms and response to clopidogrel. *N Engl J Med*. 2009;360:354-362.
12. Wang Y, Zhao X, Lin J, Li H, Johnston C, Lin Y, et al. Association between CYP2C19 loss-of-function allele status and efficacy of clopidogrel for risk reduction among patients with minor stroke or transient ischemic attack. *JAMA*. 2016;316:70-78.
13. Mirabbasi SA, Khalighi K, Wu Y, Walker S, Khalighi B, Fan W, et al. CYP2C19 genetic variation and individualized clopidogrel prescription in a cardiology clinic. *J Community*

Hosp Intern Med Perspect. 2017;7:151-156.

14. Ned RM. Genetic testing for CYP450 polymorphisms to predict response to clopidogrel: current evidence and test availability: Application: Pharmacogenomics. *PLOS Currents Evidence on Genomic Tests*. 2010 Sep 20 . Edition 1. doi: 10.1371/currents.RRN1180.
15. Doll J, Zeitler E, Becker R. Generic clopidogrel: time to substitute? *JAMA*. 2013;310:145-146.
16. Kesselheim AS, Misono AS, Lee JL, Stedman MR, Brookhart MA, Choudhry NK, et al. Clinical equivalence of generic and brand-name drugs used in cardiovascular disease: a systematic review and meta-analysis. *JAMA*. 2008;300:2514-2526.
17. Privitera MD, Welty TE, Gidal BE, Diaz FJ, Krebill R, Szafarski JP, et al. Generic-to-generic lamotrigine switches in people with epilepsy: the randomised controlled EQUI-GEN trial. *Lancet Neurol*. 2016;15:365-372.
18. Duerden MG, Hughes DA. Generic and therapeutic substitutions in the UK: are they a good thing? *Br J Clin Pharmacol*. 2010;70:335-341.
19. Kesselheim AS, Stedman MR, Bublick EJ, Gagne JJ, Misono AS, Lee JL, et al. Seizure outcomes following the use of generic versus brand-name antiepileptic drugs: a systematic review and meta-analysis. *Drugs*. 2010;70:605-621.
20. Dentali F, Donadini MP, Clark N, Crowther MA, Garcia D, Hylek E, et al. Brand name versus generic warfarin: a systematic review of the literature. *Pharmacotherapy*. 2011;31:386-393.
21. Bigdeli M, Jacobs B, Tomson G, Laing R, Ghaffar A, Dujardin B, et al. Access to medicines from a health system perspective. *Health Policy Plan*. 2013;28:692-704.