

# what's your diagnosis?

## Diagnosis: Madura foot

Nishant Nayyar, Shikha Sood, Ashwani Tomar

From the Department of Radiodiagnosis, Indira Gandhi Medical College, Shimla, Himachal, Pradesh, India

**Correspondence:** Dr. Nishant Nayyar · Department of Radiodiagnosis, Indira Gandhi Medical College, Shimla, Himachal Pradesh 171001, India · T: 09418026199 · nishant23rpgmc@gmail.com · ORCID: <http://orcid.org/0000-0003-2227-9105>

**Citation:** Nayyar N, Sood S, Tomar A. Diagnosis: Madura foot. *Ann Saudi Med* 2018; 38(6): epub DOI: 10.5144/0256-4947.2018.450a

**Received:** July 22, 2018

**Accepted:** September 13, 2018

**Published:** December 6, 2018

**Copyright:** Copyright © 2018, *Annals of Saudi Medicine*, Saudi Arabia. This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND). The details of which can be accessed at <http://creativecommons.org/licenses/by-nc-nd/4.0/>


**Funding:** None.

X-rays showed cortical erosion with a periosteal reaction in the metatarsal bones with adjacent soft tissue. Bones of the foot were osteopenic (**Figure 2**). Ultrasound of the foot showed the characteristic 'dot-in-circle' sign with multiple hypoechoic lesions on the dorsum with central fine hyperechoic areas (**Figure 3**). MRI showed a typical "dot-in-circle" sign on T2- and STIR-weighted images with a hyperintense spherical lesion with central hypointense foci (**Figure 4**). A radiological diagnosis of eumycetoma was made, which was confirmed with histopathological findings. Mycetoma pedis is a granulomatous disease of the subcutaneous tissue and deeper structures with the formation of grains containing aggregates of the causative organisms that may be discharged onto the skin surface through multiple sinuses. The process is indolent and may lead to abscess formation, osteomyelitis with severe deformity and disability if not treated expeditiously.<sup>1-4</sup> Although biopsy and culture provide a definitive diagnosis, these are time-consuming procedures; hence radiographs, ultrasound and MRI are early noninvasive diagnostic tools for early diagno-



**Figure 2.** Plain radiograph anteroposterior view showing cortical erosion in 1st and 2nd metatarsal bones with periosteal reactions in metatarsal bones. Other bones of the foot are osteopenic.

# what's your diagnosis?



**Figure 3.** Ultrasound image showing typical “dot-in-circle” sign with hypoechoic lesion with central hyperechoic fungal grains (arrows).



**Figure 4.** Sagittal T2-weighted MR image showing hyperintense lesion on dorsal aspect with central hypointense fungal grains (arrows).

sis and better outcome in these patients.<sup>5-7</sup> The typical “dot-in-circle” sign on USG helps to identify Madura foot and differentiate the causative agent, *Eumycetes*. The “dot-in-circle” consists of multiple hypoechoic nodules with central hyperechogenicity. This central hyperechogenicity is fine in actinomycetoma while large and sharp in eumycetoma.<sup>5,6</sup> The “dot-in-circle”

sign is also demonstrable on MRI where a hyperintense spherical lesion with central hypointense foci is seen. The hyperintense area represents granulomata while a low intensity hypointense center represents a fungal ball or grain.<sup>7</sup>

## REFERENCES

1. Singh H. Mycetoma in India. *Indian J Surg.* 1979;4:577-9.
2. Venugopal PL, Venugopal TL. Actinomadura madurae causing mycetomas in Madras. *Indian J Pathol Microbiol.* 1991;34:119-25.
3. Mathur DR, Joshi KR, Mathur A. An etiological and pathological study of mycetoma in western Rajasthan. *Curr Med Pract.* 1979;23:151-161.
4. Venugopal TV and Venugopal PV. Mycetoma in Madras. *Sabouraudia.* 1977;15:17-22.
5. Kumar J, Kumar A, Sethy P, Gupta S. The dot-in-circle sign of mycetoma on MRI. *Diagn Interv Radiol.* 2007;13:193-5.
6. Lee Parker BM, Singh D, Biz C. The dot-in-circle sign in Madura foot. *J Foot Ankle Surg.* 2009;48:690.
7. Cherian RS, Betty M, Manipadam MT, Cherian VM, Poonnoose PM, Oommen AT, et al. The ‘dot-in-circle’ sign—a characteristic MRI finding in mycetoma foot: A report of three cases. *Br J Radiol.* 2009;82:662-5.